




PAHIATUA	Woodville	5,000	0	10,000	10,000	30,000	5,000
	Kumeroa	40,000	0	15,000	10,000	10,000	5,000
	Ballance	25,000	0	30,000	0	5,000	0
	Mangatainoka	10,000	0	15,000	3,000	25,000	0
	Kaitawa	0	20,000				15,000
	Hamua	5,000		5,000	10,000		
	South Pahiatua		0		20,000	15,000	5,000
Eketahuna					25,000	0	
TAUMARUNUI	Ongarue	5,000					
	Western Bay	10,000					
	Matiere			20,000	55,000	40,000	20,000
	Ohura	20,000	20,000	20,000			5,000
	Burnand Road	95,000	925,000	160,000	15,000	50,000	
	Kirukau		10,000		30,000		
LEVIN	Waitare	5,000	5,000	5,000	15,000	15,000	20,000
	Ihakara	15,000	5,000		25,000	25,000	
	Shannon	15,000	10,000	15,000	40,000	40,000	45,000
	Foxton	5,000	5,000	5,000	45,000	50,000	50,000
	Moutoa	5,000	5,000	5,000	75,000	80,000	75,000
	Levin	0	0	10,000	0	5,000	20,000
	Ohau	0	10,000	5,000	5,000	0	0
	Manakau	0	0	0	5,000	0	15,000
	Otaki	0	5,000	5,000	0	0	15,000
	Te Horo	5,000	25,000	0	25,000	0	0

Spore counts can vary greatly between properties and even between paddocks. We encourage farmers to bring in grass samples to get a more accurate idea of the facial eczema risk on their own property. Contact your local Totally Vets, Tararua Veterinary Services or Levin & Horowhenua Veterinary Centre branch if you would like more information on bringing in your own samples for testing.

### FACIAL ECZEMA MONITOR SITES BY BRANCH



### Totally Vets Taumarunui

